

Ionic Liquid - Based Green Solvents for Sustainable Organic Synthesis

Dr. Manisha Singhal¹ and Dr. Shashi Bala²

¹Associate Professor, Department of Chemistry, Raghunath Girls' Postgraduate College, Meerut, Uttar Pradesh, INDIA

²Associate Professor, Department of Zoology, Raghunath Girls' Postgraduate College, Meerut, Uttar Pradesh, INDIA

¹Corresponding Author: manishasinghal2030@gmail.com



www.jrasb.com || Vol. 5 No. 2 (2026): April Issue

Received: 08-04-2026

Revised: 15-04-2026

Accepted: 19-04-2026

ABSTRACT

Ionic liquids (ILs) have emerged as promising green solvents for sustainable organic synthesis because of their negligible vapor pressure, high thermal stability, tunable polarity, and ability to function simultaneously as solvents and catalysts. In recent years, research—including significant contributions from Indian chemists—has focused on replacing volatile organic compounds with ionic liquid systems to achieve safer, energy-efficient, and environmentally benign synthetic pathways. This research paper reviews ionic liquid-based green solvent systems used in sustainable organic synthesis, covering their structural principles, physicochemical properties, synthesis approaches, reaction applications, recyclability, and environmental considerations. Methodologies discussed include solvent design, catalytic reaction optimization, and performance evaluation through yield, selectivity, and recyclability studies. Results from recent literature show that IL-based systems significantly improve reaction efficiency, reduce waste generation, and support multicomponent and heterocyclic synthesis. The discussion highlights challenges such as toxicity assessment, cost, and large-scale implementation while outlining future directions in bio-based ionic liquids and hybrid green solvent systems.

Keywords- Ionic Liquids; Green Solvents; Sustainable Organic Synthesis; Green Chemistry.

I. INTRODUCTION

Ionic liquids (ILs) are salts composed entirely of ions that remain liquid at or near room temperature, offering unique physicochemical properties that distinguish them from conventional molecular solvents [1,2]. Their low vapor pressure and non-flammability make them attractive alternatives to volatile organic solvents that contribute significantly to environmental pollution. The development of green chemistry principles has encouraged chemists to redesign synthetic processes by reducing hazardous solvent usage. Since solvents account for a large proportion of chemical waste, replacing traditional solvents with environmentally benign alternatives is a central goal of sustainable organic synthesis [3]. Ionic liquids are often described as “designer solvents” because their properties can be tuned through modification of cation–anion combinations. This tunability enables optimization for specific reactions, such as enhanced selectivity,

improved solubility, and better catalytic performance [4]. Indian researchers have actively participated in exploring ionic liquids for organic transformations, including heterocyclic synthesis, multicomponent reactions, and catalytic processes. Studies from Indian universities highlight the ability of ILs to improve reaction yields while reducing environmental impact [5,6].

The green characteristics of ILs arise primarily from their negligible vapor pressure, which minimizes atmospheric emissions compared to traditional volatile solvents. Their thermal stability further allows reactions to proceed under mild or controlled high-temperature conditions without solvent degradation [7]. In organic synthesis, ionic liquids frequently act as both reaction media and catalysts, thereby reducing the need for additional reagents. This dual functionality contributes to simplified reaction systems and decreased waste generation [8]. Recent reviews have demonstrated the effectiveness of ILs in promoting click reactions,

cycloaddition reactions, esterification, and heterocycle synthesis, with improved reaction efficiency and easier product separation [9]. Another important advantage is recyclability. Many ionic liquid systems can be recovered and reused multiple times without significant loss in catalytic performance, supporting circular and sustainable chemical processes [10].

Indian studies emphasize IL-assisted synthesis of medicinally important heterocycles, where reaction rates and selectivity are enhanced due to improved solvation of reactants and stabilization of intermediates [6]. Integration of ionic liquids with other green technologies, such as metal–organic frameworks and catalytic systems, represents an emerging research direction aimed at maximizing sustainability benefits [11]. Despite their advantages, concerns regarding toxicity, biodegradability, and energy cost of synthesis have prompted researchers to explore safer and bio-derived ionic liquids as next-generation green solvents [12]. Overall, ionic liquid–based solvent systems represent a significant shift toward environmentally responsible synthetic chemistry and have become a central topic within sustainable organic synthesis research up to 2025 [4,9].

II. METHODOLOGY

The design of ionic liquids begins with selecting appropriate cations and anions to achieve desired physicochemical properties such as polarity, viscosity, and hydrophobicity [4]. Imidazolium-, pyridinium-, and ammonium-based ionic liquids are commonly studied in sustainable synthesis applications.

Synthesis of ionic liquids generally involves quaternization reactions followed by anion exchange to produce tailored solvent systems suitable for specific reactions [7]. Purification and drying of ionic liquids are crucial steps because impurities or residual water may significantly affect catalytic performance and reaction outcomes. Characterization techniques including NMR spectroscopy, FTIR, and thermal analysis are used to confirm chemical structure and evaluate thermal stability of prepared ionic liquids [1]. Reaction performance studies typically compare ionic liquid systems against conventional organic solvents by monitoring yield, reaction time, and selectivity [8].

Many studies employ ionic liquids as catalytic media in multicomponent reactions where multiple

reactants combine in a single vessel, improving atom economy and process efficiency [6]. Recyclability testing involves separating reaction products from ionic liquids via extraction or phase separation, followed by reuse of the ionic liquid in repeated cycles [10]. Energy efficiency evaluation is performed by monitoring reaction temperature and time reduction compared with traditional solvent systems. Several studies combine ionic liquids with metal catalysts or solid supports to enhance catalytic activity and facilitate product separation [11]. Environmental evaluation includes qualitative analysis of toxicity, biodegradability, and lifecycle considerations associated with ionic liquid synthesis and disposal [12]. Mechanistic studies are conducted to understand how ionic liquids influence transition states, reaction intermediates, and catalytic pathways.

Finally, scalability assessments explore whether ionic liquid-mediated reactions can be translated from laboratory-scale synthesis to pilot-scale processes while maintaining sustainability advantages [3].

III. RESULTS

Studies up to 2025 demonstrate that ionic liquids significantly enhance yields in organic synthesis compared with traditional solvents due to improved solvation and catalytic stabilization effects. Ionic liquid-based systems show improved selectivity in cycloaddition and heterocyclic synthesis, reducing side-product formation and simplifying purification steps [9]. Reaction times are often shortened when ionic liquids are used, leading to lower energy consumption and more efficient processes. Multicomponent reactions performed in ionic liquids display higher atom economy and reduced solvent waste, aligning strongly with green chemistry principles [6]. Recyclability studies indicate that many ionic liquids maintain catalytic activity over multiple reaction cycles, supporting cost and sustainability advantages [10]. Hybrid systems combining ionic liquids with catalysts or porous materials show synergistic effects, further enhancing efficiency and selectivity [11]. However, some studies highlight that not all ionic liquids are inherently “green,” emphasizing the need for toxicity screening and lifecycle analysis [12]. Overall, literature indicates that ionic liquids provide a versatile platform for sustainable synthesis but require careful selection and design for environmentally safe application.

Table 1: Common Ionic Liquids Used in Sustainable Organic Synthesis

Ionic Liquid Type	Typical Application	Key Advantage
Imidazolium-based ILs	Cycloaddition reactions	High catalytic efficiency
Ammonium-based ILs	Multicomponent synthesis	Mild reaction conditions
Pyridinium-based ILs	Heterocyclic synthesis	Enhanced selectivity
Functionalized ILs	Catalytic transformations	Dual solvent–catalyst role

Table 2: Observed Sustainability Benefits

Parameter	Observation
Vapor pressure	Extremely low emissions
Reaction efficiency	Higher yields and selectivity
Reusability	Multiple catalytic cycles
Waste generation	Reduced solvent waste
Energy demand	Lower reaction temperature/time

IV. DISCUSSION AND CONCLUSION

Ionic liquid-based green solvents have transformed sustainable organic synthesis by offering safer alternatives to volatile organic solvents. Their ability to function as both solvent and catalyst contributes significantly to process intensification. Recent research, including Indian contributions, demonstrates that ionic liquids enable high-yield synthesis of complex organic molecules under greener conditions with reduced waste generation. Recyclability remains one of the strongest advantages of ionic liquids; however, large-scale implementation requires improved recovery methods and economic viability. Challenges remain regarding toxicity evaluation and environmental impact of ionic liquid production, emphasizing the importance of developing bio-based and biodegradable alternatives. Future research is expected to focus on hybrid solvent systems, computational solvent design, and industry-oriented scale-up strategies that maximize sustainability.

REFERENCES

[1] Phukan, P., & Sarma, D. (2021). *Synthesis of Medicinally Relevant Scaffolds—Triazoles and Pyrazoles in Green Solvent Ionic Liquids*. *Current Organic Chemistry*, 25(13), 1457–1477.

[2] Welton, T. (1999). *Room-Temperature Ionic Liquids: Solvents for Synthesis and Catalysis*. *Chemical Reviews*, 99(8), 2071–2084.

[3] Byrne, F. P., Jin, S., Paggiola, G., et al. (2016). *Tools and Techniques for Solvent Selection: Green Solvent Selection Guides*. Sustainable Chemical Processes, 4, 7.

[4] Hayes, R., Warr, G. G., & Atkin, R. (2015). *Structure and Nanostructure in Ionic Liquids*. *Chemical Reviews*, 115(13), 6357–6426.

[5] Dinodia, M. (2022). *Ionic liquids: Environment-friendly greener solvents for organic synthesis*. *Current Organic Synthesis*, 19(4), 543–557.

[6] Clarke, C. J., Tu, W.-C., Levers, O., Bröhl, A., & Hallett, J. P. (2018). *Green and Sustainable Solvents in Chemical Processes*. *Chemical Reviews*, 118(2), 747–800.

[7] Tzani, A., Kourgiantaki, A., & Detsi, A. (2022). *Current Trends in Green Solvents: Biocompatible Ionic Liquids and Deep Eutectic Solvents*. *Crystals*, 12(12), 1776.

[8] Kumar, A., et al. (2024). *Ionic liquids for the green synthesis of 1,2,3-triazoles*. *Green Chemistry*.

[9] Dwivedi, J., et al. (2025). *Catalytic Application of Ionic Liquids for the Green Synthesis of Aromatic Five-Membered Nitrogen Heterocycles*. *Catalysts*, 15(10), Article 931.

[10] Ranke, J., Stolte, S., Störmann, R., Arning, J., & Jastorff, B. (2007). *Design of Sustainable Chemical Products—The Example of Ionic Liquids*. *Chemical Reviews*, 107(6), 2183–2206.

[11] Zhu, M., et al. (2025). *Ionic Liquid Catalysis in Synthesizing Pyrazoles and Triazoles*. *Sustainable Chemistry for the Environment*.

[12] Anastas, P. T., & Warner, J. C. (1998). *Green Chemistry: Theory and Practice*. Oxford University Press.